



Formax Medicated ROL

For control of fecal flies in the manure of treated cattle. Prevents development of face flies, horn flies, house flies and stable flies in the manure of treated cattle.

Guaranteed Analysis

Calcium (CA)	(Min)	15.00	(Max)	17.00 %
Magnesium (Mg)			(Min)	2 %
Phosphorus (P)			(Min)	2.00 %
Potassium (K)			(Min)	0.25 %
Salt	(Min)	0.22	(Max)	21.00 %
Cobalt			(Min)	5 ppm
Copper (Cu)			(Min)	200 ppm
Total Selenium			(Min)	26.00 ppm
Iodine			(Min)	48 ppm
Iron (Fe)			(Min)	5,700 ppm
Manganese (MN)			(Min)	850 ppm
Zinc (Zn)			(Min)	800 ppm
Vitamin A			(Min)	50 KIU/lb
Vitamin D			(Min)	13 KIU/lb
Vitamin E			(Min)	13 IU/lb
Tetrachorvinphos			(Min)	4,500 g/ton

List of Ingredients

Calcium Carbonate, Salt, Processed Grain By-Products, Monocalcium Phosphate, Rabon 7.76%, Roughage Products, Magnesium Oxide, Molasses Products, Mineral Oil, Iron Oxide, Manganese Sulfate, Zinc Sulfate, Potassium Chloride, Artificial Flavoring, Copper Sulfate, Sodium Selenite, Calcium Iodate, Vitamin D3 Supplement, Cobalt Carbonate, Vitamin E Supplement, Vitamin A Supplement, Propionic Acid, Acetic Acid.

SS003711 Feeding Directions

See Back of label for feeding directions.

WARNING: This product, which contains added copper, should not be fed to sheep or any species with a low tolerance to supplemental copper.

CAUTION: Follow label directions.

Consumption of selenium should not exceed 3 mg per head per day.

Manufactured for:

ALABAMA FARMERS COOPERATIVE

Decatur, Alabama 35609

NET WEIGHT: 50 LBS. (22.7 KG)

15/07/11



PRECAUTIONARY STATEMENTS
Hazardous to Humans and Domestic animals.

CAUTION:

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing.

User Safety Guidelines

Follow all label instructions.

- Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 to 2 glasses of water. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON CLOTHING OR SKIN: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 – 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Contains an organophosphate that inhibits cholinesterase.

Have the product container or label with you when you call a poison control center or doctor, or going for treatment. You may also contact 1-800-821-7467 for emergency medical treatment information.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Formax Medicated ROL is formulated to be fed continuously on a free choice basis to cattle to provide minerals, vitamins and fly control daily. Feed supply and feeder locations should be adequate to assure consumption in accordance with the intake levels stated below.

CATTLE WEIGHT	Formax Medicated ROL daily intake		Tetrachorvinphos Daily Intake.
	(Lb.)	(Oz.)	(Lb.)
500	2.5	0.15625	350
600	3	0.18750	420
700	3.5	0.21875	490
800	4	0.25000	560
900	4.5	0.28125	630
1000	5	0.31250	700
1100	5.5	0.34375	770
1200	6	0.37500	840

If feed consumption varies up or down from this chart, move mineral feeders further from or closer to general watering and loafing areas. If cattle still fail to consume the correct amount, discontinue feeding this free choice supplement. Feed Formax Medicated ROL as the only source of salt, other minerals and vitamins.