

CO-OP GAME BIRD STARTER-GROWER

MEDICATED

ITEM NUMBER 270320

MEDICATED:

For the control of necrotic enteritis caused or complicated by *Clostridium* spp. or other organisms susceptible to Bacitracin

Active Drug Ingredient:

Bacitracin Methylene Disalicylate 200GM/TN

GUARANTEED ANALYSIS

CRUDE PROTEIN, not less than	27.0%
CRUDE FAT, not less than	3.0%
CRUDE FIBER, not more than	5.0%
CALCIUM, not less than	1.2%
CALCIUM, not more than	1.3%
PHOSPHOROUS, not less than	.9%

Ingredients

Grain Products, Plant Protein Products, Processed Grain By-Products, Forage Products, Calcium Carbonate, Roughage Products, Poultry Byproducts, Dicalcium Phosphate, Salt, Vitamin A Supplement, Vitamin D Supplement, Vitamin E Supplement, Vitamin B-12 Supplement, Menadione Sodium Bisulfate Complex, Choline Chloride, Riboflavin Supplement, Thiamine Mononitrate, Niacin, Folic Acid, Biotin, Pyridoxine Hydrochloride, Ferrous Sulfate, Copper Sulfate, Zinc Oxide, Manganous Oxide, Sodium Selenite, and DL Methionine

Manufactured By

ALABAMA FARMERS COOPERATIVE

Decatur, Alabama 35601

Feeding Directions:

CO-OP Game bird Medicated Starter Grower is a highly fortified poultry feed designed with the game bird baby chick in mind. This feed is also medicated to promote growth and prevent coccidiosis, the leading cause of chick mortality. This 27% protein feed contains extra vitamins and minerals to reduce stress, and sickness in young birds. The minerals and vitamins will increase their immune system to fight of disease and sickness, while the extra protein will promote growth and meat production. The extra calcium will also aid to the development of bone as the bird develops. This feed also contains high percentages of the amino acids Lysine, and Methionine, which when deficient is the amino acids that most

limit growth in poultry. This feed is recommended for a sole ration for chicks until they reach 20 weeks of age. At 20 weeks of age it is recommended to slowly switch birds to CO-OP Game Bird Booster. Do not feed this product as a laying feed

Important Management Practices:

Always provide a source of fresh, clean water, Do not allow feed to become wet or moldy. Always keep feeders and waterers clean. Do not make sudden changes in the diets of game birds. When changing feeds, allow a 7 to 10 day transition period.