

SFGD CATTLE WORMER PELLET

Item number 234120

MEDICATED

For the removal and control of gastrointestinal worms: Large Roundworms (*Ascaris suum*, Nodular Worms (*Hyostrongylus rubidus*), Whipworm (*trichuris suis*) Kidney Worms (Mature and Immature *Stephanurus dentatus*), and Lungworms (*Metastrongylus apri*)

ACTIVE DRUG INGREDIENT

Fenbendazole

567MG/LB

Caution:

Do not use within 13 days of slaughter. Do not use in dairy cattle of breeding age.

GUARANTEED ANALYSIS:

Crude Protein, (not less than)	20.00%
Crude Fat, (not less than)	2.00%
Crude Fiber, (not more than)	15.00%

INGREDIENTS:

Grain Products, Plant Protein Products, Processed Grain By-Products, Calcium Carbonate, Roughage Products, Forage Products, Molasses Products, Dicalcium Phosphate, Salt, Fenbendazole, Magnesium Oxide, Vitamin A Supplement, Vitamin D3 Supplement, Vitamin E Supplement, , Ferrous Sulfate, Copper Sulfate, Zinc Oxide, Maganous Oxide, Sodium Selenite, Zinc Sulfate, Maganous Sulfate and Cobalt Carbonate

Manufactured By

ALABAMA FARMERS COOPERATIVE, INC.

Decatur, Alabama 35601

Feeding Directions:

CO-OP Safeguard Worming Pellets is a popular 20% protein product designed to deworm pasture cattle. Parasites cost producers more money each year than any other non feed item. The active worming ingredient in this product is effective in the control of Stomach Worms, Intestinal Worms, and Lungworms. This product will not control external parasites such as flies and lice. This feed also contains cottonseed meal, soybean meal, alfalfa meal, and corn, which cows will readily consume. Feed this product at the rate of 1lb per 250 pounds of body weight. Safeguard Worming Pellets is a very economical and labor free method of cattle worming. Feed as sole ration for one day, making sure all cattle consume product at recommended rate.

Important Management Practices

Provide a source of fresh clean water and free choice mineral at all times. When making any change in an animal's diet, plan a 7 to 10 day transition period. Provide ample feeder space of 18 inches per animal. Feeders should protect feed from becoming wet and moldy. Poorly managed feeding programs can increase the possibility of feeding disorders. You should contact an Animal Nutritionist when using this product in a way other than recommended.