



01S03315

# Formax Fly Control IGR Breeder Mineral

PREVENTS ADULT HORN FLY EMERGENCE FROM MANURE OF TREATED CATTLE

A nutritional supplement containing S-Methoprene\* Insect Growth Regulator for continuous feeding to cattle through the horn fly season.

**ACTIVE INGREDIENT:**

S-Methoprene (CAS #65733-16-6).....0.01%

**OTHER INGREDIENTS:** .....99.99%

**TOTAL** .....100.00%

\*As the Insect Growth Regulator ALTOSID®

**GUARANTEED ANALYSIS**

Calcium (Ca), min.....18.30%	Copper (Cu), min.....1,800 ppm
Calcium (Ca), max.....21.30%	Manganese (Mn), min.....5,400 ppm
Phosphorus (P), min.....4.00%	Selenium (Se), min.....26.00 ppm
Salt (NaCl), min.....13.50%	Zinc (Zn), min.....5,400 ppm
Salt (NaCl), max.....16.50%	Vitamin A, min.....200,000 IU/lb
Magnesium (Mg), min.....2.00%	Vitamin D3, min.....20,000 IU/lb
Potassium (K), min.....1.00%	Vitamin E, min.....100 IU/lb

**FEED INGREDIENTS**

Calcium Carbonate, Monocalcium Phosphate, Salt, Roughage Products, Magnesium Oxide, Processed Grain By-Products, Molasses Products, Potassium Chloride, Vegetable Oil, Calcium Stearate, Manganese Sulfate, Copper Sulfate, Iron Oxide, Zinc Sulfate, Selenium Yeast, Artificial Flavoring, Sodium Selenite, Vitamin D<sup>3</sup> Supplement, Ethylenediamine Dihydroiodide, Cobalt Carbonate, Vitamin A Acetate, Vitamin E Supplement, Copper Methionine Hydroxy Analogue Chelate, Zinc Methionine Hydroxy Analogue Chelate, Manganese Methionine Hydroxy Analogue Chelate.

**KEEP OUT OF REACH OF CHILDREN**  
**WARNING**

**See additional Precautionary Statements**

EPA Reg. No. 2724-494-88091

EPA Est No. 88091-TN-001

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS - WARNING:** This product contains high levels of copper. Consumption of this product by sheep or goats may result in copper toxicity. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses). Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

**FIRST AID:**

If in Eyes:	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
-------------	---

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 for emergency medical treatment information.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

## FEEDING AND MANAGEMENT INSTRUCTIONS

When used as directed, Formax Fly Control IGR Breeder Mineral will provide sufficient S-methoprene insect growth regulator to prevent the emergence of adult horn flies from manure of treated cattle. Existing adult horn flies will not be affected. Start feeding before horn flies appear and continue use until cold weather marks the end of the horn fly season. Allow free choice to cattle. Cattle should consume an average of 0.4 ounces of Formax Fly Control IGR Breeder Mineral per 100 lbs. of body weight per day, which is equivalent to 4 ounces per day for a 1,000 pound animal.

1. Assure cattle are not starved for nutrients in Formax Fly Control IGR Breeder Mineral by providing them prior to feeding Formax Fly Control IGR Breeder Mineral.
2. Offer cattle 50 pounds of Formax Fly Control IGR Breeder Mineral for every 15 head of cattle.
3. Remove all other self limiting nutritional supplements.
4. Locate Formax Fly Control IGR Breeder Mineral where cattle congregate (watering, loafing, shade areas).
5. Replenish Formax Fly Control IGR Breeder Mineral on a regular basis as needed.
6. If intake of Formax Fly Control IGR Breeder Mineral is below 0.4 ounces per 100 lbs. animal body weight per day, increase the number of feeding locations and/or change location(s) to areas more frequented by cattle.
7. If intake of Formax Fly Control IGR Breeder Mineral is above 0.4 ounces per 100 lbs. animal body weight per day, reduce the number of feeding locations and/or change location(s) to areas less frequented by cattle.
8. Introduction of this product after adult horn fly infestation is established will require treatment of cattle with adulticides if elimination of the adult fly population is desired.

**STORAGE AND DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal.

**Storage:** Store in a cool place.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Completely empty bag, then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For information, call 1-800-552-9620

ALTOSID® Insect Growth Regulator is a registered trademark of Wellmark International.

Made in USA

Manufactured by:  
**Gold River Feed Products**  
783 Eagle Blvd  
Shelbyville, TN 37160  
NET WEIGHT: 50 LBS. (22.7 KG)