



# Formax Goat Mineral R MEDICATED

For the prevention of coccidiosis caused by *Eimeria crandallis*, *E. christensenii*, and *E. ninakohlyakimovae*.

### ACTIVE DRUG INGREDIENT

Monensin Sodium ..... 1460 g/ton

### Guaranteed Analysis

Calcium	(Min)	15.00 %	Copper	(Min)	1,000 ppm
Calcium	(Max)	18.00 %	Iodine	(Min)	80 ppm
Phosphorus	(Min)	8.00 %	Manganese	(Min)	3,000 ppm
Salt	(Min)	18.00 %	Selenium	(Min)	26 ppm
Salt	(Max)	22.00 %	Zinc	(Min)	3,000 ppm
Magnesium	(Min)	2.00 %	Vitamin A	(Min)	200,000 IU/lb
Potassium	(Min)	1.00 %	Vitamin D	(Min)	20,000 IU/lb
Cobalt	(Min)	30 ppm	Vitamin E	(Min)	20 IU/lb

### List of Ingredients

Calcium Carbonate, Salt, Monocalcium Phosphate, Processed Grain By-Products, Roughage Products, Magnesium Oxide, Molasses Products, Potassium Chloride, Ammonium Chloride, Vegetable Oil, Iron Oxide, Manganese Sulfate, Zinc Sulfate, Copper Sulfate, Artificial Flavoring, Sodium Selenite, Vitamin D3 Supplement, Calcium Iodate, Cobalt Carbonate, Iron Carbonate, Manganous Oxide, Zinc Oxide, Ferrous Sulfate, Dried Extracted *Aspergillus niger* Fermentation Solubles, Choline Chloride, Niacin Supplement, Vitamin E Supplement, Vitamin A Supplement, Calcium Pantothenate, Riboflavin Supplement, Silicon Dioxide, Amorphous Silicon Dioxide, Phosphoric Acid, Citric Acid (a preservative), Disodium EDTA Butylated Hydroxyanisole (a preservative), Menadione Sodium Bisulfate Complex, Biotin, Thiamine Mononitrate, Folic Acid, Pyridoxine Hydrochloride, Vitamin B12 Supplement, Propionic Acid, Acetic Acid, Copper Methionine Hydroxy Analogue Chelate, Zinc Methionine Hydroxy Analogue Chelate, Manganese Methionine Hydroxy Analogue Chelate.

### Feeding Directions

#### TYPE B MEDICATED ARTICLE

Mix Formax Goat Mineral R thoroughly with non-medicated feed to provide 20g Monensin per ton of complete feed. Thoroughly mix 27.60 lb. of Formax Goat Mineral R with 1972.4 lb. of non-medicated complete feed. Provide clean fresh water at all times.

CAUTION: USE AS DIRECTED

THIS FEED SHOULD BE USED ONLY IN ACCORDANCE WITH THE RECOMMENDATIONS ON THIS FEED TAG

CAUTION: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by equines has been fatal. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal in cattle and could be fatal in goats. Monensin medicated feed is safe for use in goats only. Consumption by unapproved species may result in toxic reactions.

LIMITATIONS: Feed continuously. Do not feed to lactating goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the level of Monensin recommended in the feeding direction, as reduced average daily gains may result.

CAUTION: Follow label directions.

Consumption of selenium should not exceed 3 mg per head per day. Feeding 4.0 oz. of Formax Goat Mineral R will provide 3mg Selenium per head daily.

This product was manufactured in a facility that does not handle or store products containing animal proteins prohibited in ruminant feed as described in 21 CFR section 589.2000.

Rumensin® is a registered trademark of Elanco Animal Health.

Manufactured for:

**ALABAMA FARMERS COOPERATIVE**

**Decatur, Alabama 35609**

**NET WEIGHT: 50 LBS. (22.7 KG)**

Net Weight 50 lb. or Bulk Invoice

