

# HORIZON SENIOR

263040

*Pelleted*  
**FOR MATURE HORSES**

---

## Guaranteed Analysis

|               |       |       |       |
|---------------|-------|-------|-------|
| Crude Protein | (Min) | 14.50 | %     |
| Crude Fat     | (Min) | 7.00  | %     |
| Crude Fiber   | (Max) | 18.00 | %     |
| Calcium       | (Min) | 0.75  | %     |
| Calcium       | (Max) | 1.75  | %     |
| Phosphorus    | (Min) | 0.60  | %     |
| Vitamin A     | (Min) | 8,000 | IU/lb |
| Vitamin D     | (Min) | 500   | IU/lb |
| Vitamin E     | (Min) | 100   | IU/lb |

---

## Ingredients

Processed Grain By-Products, Forage Products, Plant Protein Products, Cane Molasses, Vegetable Oil, Propionic Acid, SOY OIL, Molasses Products, Animal Fat, Calcium Carbonate, Animal Protein Products, Grain Products, Dicalcium Phosphate, Salt, Saccharomyces Yeast Culture, KAOLIN, Monosodium Phosphate, Potassium Sulfate, Magnesium Sulfate, GHP 2, MINTREX, Ferrous Sulfate, Zinc Sulfate, Manganous Sulfate, Copper Sulfate, Cobalt Carbonate, Calcium Iodate, Sodium Selenite, Mineral Oil, Iron Oxide, Vitamin E Supplement, Biotin, Riboflavin, Thiamine Mononitrate, Calcium Pantothenate, Vitamin A Supplement, Niacin Supplement, Folic Acid, Vitamin D 3 Supplement, Vitamin B12 Supplement, Roughage Products, DEXTROSE, Mold Inhibitor, Anise, MTB100 55#, EQUINE BIO FORM., Lysine, Zinc Methionine Complex, Manganese Methionine Complex, Copper Lysine Complex, Cobalt Gluconate, Manganous Oxide, Zinc Oxide, Sodium Selenite, Manganese Sulfate, Calcium Iodate and BIOTIN 100.

## Feeding Directions

Feed to horses at the rate of .5 to 1.5 lbs. per 100 lbs. bodyweight depending on the quality and quantity of other feedstuffs available and the intended use of the horse. Do not feed free choice to any animal. Provide good quality hay or pasture.

## CAUTION

***Do not feed to sheep due to copper levels in this feed.***

IMPORTANT MANAGEMENT PRACTICES: Always provide a fresh source of clean water, salt and free choice minerals at all times. When making any change in an animal's diet, plan a 7 to 10 day transition period. Do not allow feed to stay in feeder for a period of longer than 3 days. When feeding large amounts of this feed (over 5 pounds per day) split the feeding amount into two different feedings to reduce the incidence of colic. Poorly managed feeding programs can lead to nutritional disorders.

Manufactured by  
ALABAMA FARMERS COOPERATIVE INC.

Decatur, Alabama 35601  
Net Wt. 50 lbs. (22.67 kg)

01/15/2016



7 16374 63040 5