


Triple Crown Low Starch Formula

A low carb and corn, molasses and alfalfa meal free feed for hard keepers

Not all horses are easy keepers. Some require additional calories for higher levels of performance or elevated levels of metabolism. Since Triple Crown Low Starch has lower carbs than hay and pasture, it can be used as a complete feed to replace all or part of (the forage) portion of the diet, which typically has unknown carbohydrate and starch levels. This formula contains no corn, molasses or alfalfa meal—ingredients many horse owners believe cause erratic behavior. By utilizing good quality fiber sources and added fat, including rice bran and flaxseed, we can replace a number of the calories that are sacrificed by keeping carbs low. *Available in 50 lb. bags.*

Guaranteed Analysis

| | |
|--|------------------------|
| Crude Protein (min.) | 13.00% |
| Lysine (min.) | 0.70% |
| Methionine (min.) | 0.20% |
| Threonine (min.) | 0.35% |
| Crude Fat (min.) | 6.00% |
| Crude Fiber (max.) | 18.00% |
| Calcium (min.) | 0.75% |
| Calcium (max.) | 1.25% |
| Phosphorus (min.) | 0.60% |
| Magnesium (min.) | 0.50% |
| Iron (min.) | 170 ppm |
| Potassium (min.) | 0.75% |
| Selenium (min.) | 0.60 ppm |
| Zinc (min.) | 170 ppm |
| Manganese (min.) | 100 ppm |
| Copper (min.) | 50 ppm |
| Vitamin A (min.) | 6,000 IU/lb. |
| Vitamin D (min.) | 500 IU/lb. |
| Vitamin E (min.) | 200 IU/lb. |
| Ascorbic Acid (min.) | 45 mg/lb. |
| Biotin (min.) | 0.30 mg/lb. |
| Lactobacillus Acidophilus | 908 million CFU/lb. |
| Fermentation Product (min.) | |
| Saccharomyces Cerevisiae | 1,816 million CFU/lb. |
| Yeast Culture (min.) | |
| Cellulase (Trichoderma Longibrachiatum | 110 CMC-ase Units/lb. |
| Fermentation Extract) (min.) | |
| Protease (Bacillus Subtilis Fermentation | 0.4 Northrup Units/lb. |
| Extract) (min.) | |



TRIPLE CROWN[®]

N u t r i t i o n , I n c .

Ingredients

Wheat Middlings, Soybean Hulls, Shredded Beet Pulp, Distillers Dried Grains, Soybean Oil, Ground Limestone, Salt, Sodium Bicarbonate, Sodium Sesquicarbonate, Stabilized Rice Bran, Calcium Carbonate, Vitamin E Supplement, Yeast Culture, Hydrolyzed Yeast, Hydrated Sodium Calcium Aluminosilicate, Dried Lactobacillus Acidophilus Fermentation Product, Dried Enterococcus Faecium Fermentation Product, Anise Seed, Fenugreek Seed, Dried Trichoderma Longibrachiatum Fermentation Extract, Dried Bacillus Subtilis Fermentation Extract, Zinc Proteinate, Manganese Proteinate, Kelp Meal, Copper Proteinate, Lecithin, Magnesium Oxide, Iron Proteinate, Magnesium Proteinate, Selenium Yeast, Flaxseed, Monocalcium Phosphate, Dicalcium Phosphate, Defluorinated Phosphate, Lignin Sulfonate, L-Lysine, Ascorbic Acid (source of Vitamin C), Niacin Supplement, Biotin, Vitamin A Supplement, Thiamine Mononitrate, Beta Carotene, Calcium Pantothenate, Riboflavin Supplement, Pyridoxine Hydrochloride, Vitamin B₁₂ Supplement, Vitamin D₃ Supplement, Choline Chloride, Menadione Sodium Bisulfite Complex, (source of Vitamin K activity), Folic Acid, Brewers Dried Yeast, DL-methionine, DL-methionine Hydroxy Analog, Ferrous Sulfate, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Cobalt Sulfate, Ethylenediamine Dihydroiodide. Free from Restricted Ruminant Protein Products per Title 21, CFR 589.2000

Feeding Directions

Triple Crown Low Starch can be fed to all horses one year of age or older. Feed enough to maintain desired body condition and energy level, along with hay, and/or pasture, water and salt. If feeding less than 6 lb. per day to horses weighing 900 lb. or more, feed additional minerals and vitamins with either Triple Crown 30% or Triple Crown Lite.

While Low Starch is a complete diet and can replace all or a part of the fiber portion of your horse's diet, we strongly recommend you continue to feed long stem hay or pasture.

NOTE: Provide plenty of fresh, clean water at all times. Keep product fresh in cool, dry storage. Examine product daily. DO NOT use if old, moldy or contaminated with insects.